

Neuroscience of motivation: reward, dopamine, and driven behavior

Neurociencia de la motivación: recompensa, dopamina y conducta impulsada

Neurociência da motivação: recompensa, dopamina e comportamento dirigido

**Ícaro Soares de Carvalho
Pinheiro^{1*}**

ORCID: 0000-0002-3051-223X

Álvaro Matheus Costa Santana¹

ORCID: 0009-0003-1683-7438

Rebeca Lima Cortez Abreu¹

ORCID: 0000-0001-6027-7966

Raquel Pereira da Cruz Silva²

ORCID: 0000-0003-1034-1143

**Victor Gabriel da Costa Pimentel
de Morais¹**

ORCID: 0009-0001-2794-9545

Irlanna Thamirys Barbosa Silva¹

ORCID: 0009-0007-8775-5834

Tânia Rodrigues Furtado¹

ORCID: 0000-0002-8171-1140

Andrienny Santana da Silva³

ORCID: 0009-0004-4929-4836

Vyrna Rebeca de Carvalho Alves⁴

ORCID: 0000-0002-1641-908X

**Maria Edillayne de Assunção
Silva⁴**

ORCID: 0000-0001-7147-8157

¹Universidade Estadual do Piauí.
Piauí, Brazil.

²Centro Universitário Adventista de
Ensino do Nordeste. Bahia, Brazil.

³Faculdade do Piauí. Piauí, Brazil.

⁴Universidade Federal do Piauí. Piauí,
Brazil.

How to cite this article:

Pinheiro ISC, Santana ÁMC, Abreu RLC, Silva RPC, Morais VGCP, Silva ITB, Furtado TR, Silva AS, Alves VRC, Silva MEA. Neuroscience of motivation: reward, dopamine, and driven behavior. Glob Acad Nurs. 2026;7(Sup.1):e535. <https://dx.doi.org/10.5935/2675-5602.20200535>

*Corresponding author:

icaropinheiro@aluno.uespi.br

Submission: 01-16-2026

Approval: 03-05-2026

Abstract

Motivation for goal-directed behaviors depends on the hierarchical orchestration between brain systems of valuation and executive control. This integrative review synthesized evidence that dopamine, acting on the mesocorticolimbic axis, primarily signals reward prediction error and mobilizes the effort necessary for action, underpinning the motivational "wanting" component. This signal is integrated by corticostriatal circuits, in which the functional connectivity between the ventral striatum and the dorsolateral prefrontal cortex is essential for transforming valuation into persistent action plans capable of overcoming costs and distractions. Dysregulation of this system generates opposing psychopathologies: hypofunction and frontostriatal disconnection are associated with apathy and anhedonia in depression, while hypersensitization of the reward system to specific stimuli sustains compulsive seeking in addiction. It is concluded that adaptive motivation requires dynamic balance and effective communication within this network, offering an integrated model for the development of targeted interventions.

Descriptors: Motivation; Reward; Dopamine; Goal-Directed Behavior; Cognitive Neuroscience.

Resumen

La motivación para las conductas dirigidas a objetivos depende de la orquestación jerárquica entre los sistemas cerebrales de valoración y control ejecutivo. Esta revisión integrativa sintetizó evidencia de que la dopamina, al actuar sobre el eje mesocorticolímbico, señala principalmente el error de predicción de recompensa y moviliza el esfuerzo necesario para la acción, lo que sustenta el componente motivacional de "deseo". Esta señal se integra mediante circuitos corticoestriatales, en los que la conectividad funcional entre el estriado ventral y la corteza prefrontal dorsolateral es esencial para transformar la valoración en planes de acción persistentes capaces de superar costos y distracciones. La disregulación de este sistema genera psicopatologías opuestas: la hipofunción y la desconexión frontoestriatal se asocian con apatía y anhedonia en la depresión, mientras que la hipersensibilización del sistema de recompensa a estímulos específicos sustenta la búsqueda compulsiva en la adicción. Se concluye que la motivación adaptativa requiere un equilibrio dinámico y una comunicación eficaz dentro de esta red, lo que ofrece un modelo integrado para el desarrollo de intervenciones específicas.

Descriptores: Motivación; Recompensa; Dopamina; Conducta Dirigida a Metas; Neurociencia Cognitiva.

Resumo

A motivação para comportamentos dirigidos a objetivos depende da orquestração hierárquica entre sistemas cerebrais de valoração e controle executivo. Esta revisão integrativa sintetizou evidências de que a dopamina, atuando no eixo mesocorticolímbico, sinaliza primariamente o erro de predição de recompensa e mobiliza o esforço necessário para a ação, fundamentando o componente de "desejo" (*wanting*) motivacional. Este sinal é integrado por circuitos corticoestriatais, nos quais a conectividade funcional entre o estriado ventral e o córtex pré-frontal dorsolateral é essencial para transformar a valoração em planos de ação persistentes, capazes de superar custos e distrações. A disregulação deste sistema gera psicopatologias opostas: a hipofunção e a desconexão frontoestriatal estão associadas à apatia e anedonia na depressão, enquanto a hipersensibilização do sistema de recompensa a estímulos específicos sustenta a busca compulsiva no vício. Conclui-se que a motivação adaptativa requer o equilíbrio dinâmico e a comunicação eficaz dentro desta rede, oferecendo um modelo integrado para o desenvolvimento de intervenções direcionadas.

Descritores: Motivação; Recompensa; Dopamina; Comportamento Dirigido a Objetivos; Neurociência Cognitiva.



Introduction

Motivation is a fundamental psychological process that provides behavior with direction, intensity, and persistence, enabling organisms to satisfy homeostatic needs and pursue complex goals. In humans, these processes extend beyond immediate survival, underpinning the pursuit of knowledge, social achievement, and long-term goals and projects. Motivation is therefore central to adaptation, learning, and well-being¹. Contemporary neuroscience investigates the biological substrates of these phenomena, seeking to elucidate how specific brain circuits translate internal needs and external stimuli into coordinated, purposeful sequences of actions. Understanding this neural architecture is necessary not only to elucidate typical human behavior but also to shed light on a broad range of neuropsychiatric disorders in which motivational systems are profoundly altered².

Historically, the study of motivation has been dominated by psychological and behavioral perspectives. However, recent decades have witnessed a transformative convergence with the biological sciences, driven by the advent of brain imaging techniques and selective neural manipulation. This integration has enabled increasingly precise mapping of the neural networks involved in motivation, revealing a complex, distributed system rather than a single motivational center³. Within this context, the concept of “reward” has emerged as a cornerstone, conceptualized not merely as a pleasurable stimulus but as an outcome that guides behavior, reinforces learning, and consolidates memories. The brain’s ability to assign motivational value to stimuli and actions is therefore an essential evolutionary mechanism underlying adaptive decision-making⁴.

At the core of this valuation machinery lies the neurotransmitter dopamine, whose role beyond motor control revolutionized the field. Evidence has established that dopaminergic neurons in the ventral tegmental area (VTA) and substantia nigra pars compacta project to limbic and cortical structures, forming the central axis of the so-called mesocorticolimbic “reward system”⁵. The activity of these neurons is intrinsically associated with prediction, desire, and reinforcement learning. Contrary to the simplistic view that dopamine encodes hedonic pleasure itself (i.e., the “liking” of a reward), the current consensus holds that dopaminergic signaling is more closely associated with incentive “wanting” and the detection of discrepancies between predicted and received rewards, known as reward prediction error. This signaling is essential for directing attention and behavioral effort toward relevant stimuli and actions^{6,7}.

This dopaminergic model of reward prediction provides a mechanism for explaining how behaviors are shaped and maintained. When an action results in a reward that is better than expected, a burst of dopamine release strengthens the synaptic connections associated with that behavioral sequence, increasing the likelihood of its future repetition. Conversely, the omission of an expected reward leads to a decrease in dopaminergic activity, signaling the need to adjust the strategy. Thus, dopamine acts as a neural

teaching signal, gradually guiding the organism toward more effective behaviors for obtaining resources through a process known as reinforcement learning. This function is mediated by dopaminergic projections to the nucleus accumbens and prefrontal cortex, regions that are critical for integrating motivational, emotional, and cognitive information^{8,9}.

On this neurochemical basis, truly goal-directed behavior emerges from the dynamic interaction between brain systems that compute goal value and those responsible for planning and executing the actions required to attain it. Whereas the dopamine-rich ventral striatum is more closely involved in motivation and incentive wanting, the dorsal striatum and dorsolateral prefrontal cortex are recruited for the selection and control of specific actions, particularly when behavior requires planning, persistence in the face of obstacles, or the inhibition of competing responses¹⁰. Thus, motivation is not a unitary state but a multidimensional construct encompassing processes ranging from basic drive to the execution of complex plans, with each dimension supported by partially overlapping yet dissociable neural circuits.

Dysregulation of these motivational circuits contributes to the etiology of several clinical conditions of major public health relevance. On the one hand, dopaminergic hypofunction and anhedonia, defined as a reduced capacity to anticipate or experience pleasure, are core features of disorders such as major depressive disorder and schizophrenia with negative symptoms, contributing to apathy, loss of initiative, and severe functional impairment¹¹. On the other hand, pathological hyperactivity or sensitization of the reward system is closely associated with substance use disorders, in which drugs of abuse hijack dopaminergic learning mechanisms, producing compulsive craving and drug-seeking behavior that overrides other goals¹². These conditions illustrate the pathological extremes of a motivational continuum, highlighting the importance of precise neurochemical regulation.

Despite significant advances, fundamental questions remain unresolved and constitute the central problem addressed by this article. The first concerns the temporal and functional integration of the different components of motivation: “How are dopaminergic valuation and prediction signals originating in subcortical regions translated into structured action plans in the prefrontal cortex?” Second, ongoing debate concerns the specificity of neural circuits underlying different types of motivation, including intrinsic versus extrinsic motivation and approach-oriented versus threat-avoidance motivation, as well as how these circuits interact with higher-order cognitive processes such as decision-making and self-regulation¹³. Finally, translating this basic knowledge into more effective clinical interventions remains challenging, requiring a more precise understanding of how to selectively modulate specific components of the motivational system without disrupting its homeostatic balance.

The hypothesis guiding this review is that motivated, goal-directed behavior results from hierarchical coordination across multiple levels of processing, in which



dopamine acts as a critical modulator that tunes the sensitivity of corticostriatal networks to reward and effort signals. It is proposed that dysfunctions in this system do not simply reflect a global dopamine deficit or excess but rather dysregulation in the precision of reward prediction error signaling and/or disconnection between valuation circuits (nucleus accumbens) and circuits responsible for action control and planning (prefrontal cortex and dorsal striatum). This integrative perspective may help explain the diversity of observed clinical phenotypes.

Against this background, the objective of this review is to synthesize contemporary neuroscientific evidence on the neural architecture of motivational processes, focusing specifically on the role of dopamine-mediated reward systems in the emergence and direction of goal-directed behavior. The analysis will focus on human neuroimaging data, neurophysiological studies in animal models, and pharmacological research, which together provide an updated model of how the brain transforms motivation into purposeful action.


Methodology

This integrative literature review was designed to synthesize and analyze neuroscientific evidence on motivational processes. This method allows the inclusion of studies with different methodological designs, enabling a comprehensive and up-to-date understanding of the topic and supporting both evidence-based practice and the identification of gaps for future research¹⁴. An integrative review was selected because of its ability to combine theoretical and empirical data from multiple sources,

The methodological process was structured into six sequential stages, as proposed by Whittemore and Knafl¹⁶: (1) identification of the problem and definition of the research question; (2) systematic literature search; (3) critical appraisal of the collected data; (4) data analysis and synthesis; (5) presentation of the results; and (6) conclusion and validation. The guiding research question was: "What is the contemporary evidence regarding the neural architecture of motivational processes, with a focus on the role of dopamine-mediated reward systems in the emergence and direction of goal-directed behavior?".

The search strategy was systematically implemented between April 15 and 30, 2024, in the electronic databases PubMed, the Scientific Electronic Library Online Brazil (SciELO Brazil), and the Virtual Health Library (VHL), which aggregates several sources, including LILACS. Controlled vocabulary terms from Medical Subject Headings (MeSH) and Health Sciences Descriptors (DeCS), along with their Portuguese and English equivalents, were used in combination with Boolean operators. The search strategy was as follows: ("Motivation"[Mesh] OR "Reward"[Mesh]) AND ("Dopamine"[Mesh] OR "Dopaminergic Neurons"[Mesh]) AND ("Goal"[Mesh] OR "Goal-Directed Behavior"[Mesh] OR "Decision Making"[Mesh]). Equivalent semantic and syntactic adaptations were applied. A publication date filter covering January 2020 through April 2024 was used to ensure the currency of the synthesis.

Figure 1. Search and study selection. Rio de Janeiro, RJ, Brazil, 2020–2024



The eligibility criteria were precisely defined. The inclusion criteria were: (I) original articles, including experimental, neuroimaging, and electrophysiological studies, and review articles, including systematic, scoping,

and narrative reviews, that directly addressed the neural systems underlying motivation, reward, and/or dopamine in relation to goal-directed behavior; (II) studies conducted in humans or animal models; (III) articles published in

Portuguese, English, or Spanish; (IV) articles with full text available online; and (V) articles indexed and published within the defined period (2020–2024). The exclusion criteria were: (I) editorials, letters, conference abstracts, dissertations, and theses; (II) studies whose primary focus was not the neurobiology of motivation, such as exclusively pharmacokinetic studies or genetic studies without behavioral correlations; and (III) articles with inadequately described methodologies or potentially serious methodological biases identified during critical appraisal.

The study selection process followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA 2020) guidelines to ensure transparency and reproducibility¹⁷. The PRISMA flow diagram details the process from identification to final inclusion. Initially, the records identified in the databases ($n = 2,180$) were imported into EndNote Web reference management software, where duplicate records ($n = 394$) were removed through automated and manual procedures. During the screening phase, two independent reviewers assessed the titles and abstracts of the remaining records ($n = 1,786$), applying the eligibility criteria. The articles considered potentially relevant ($n = 263$) were retrieved in full text and assessed for eligibility. At this stage, the same two reviewers independently read the full-text articles, excluding those that did not meet the criteria and documenting the reasons for exclusion. Disagreements at any stage were resolved by consensus or, when necessary, through arbitration by a third researcher. The studies that met the eligibility criteria ($n = 11$) were ultimately included in the review.

For data extraction and organization, a standardized instrument was developed using Microsoft Excel. The instrument included fields for authors, year of publication, country of origin, study design, sample/participants, objectives, principal methods (e.g., type of neuroimaging, behavioral paradigm, and intervention), findings relevant to the research question, and main conclusions. This spreadsheet provided a synoptic overview to support the analysis.

Qualitative data analysis was conducted using thematic analysis, following the approach proposed by Braun and Clarke¹⁸, which comprises six phases: (1) familiarization with the data through initial and repeated reading of the included articles; (2) generation of initial codes by identifying relevant units of meaning; (3) searching for themes by grouping codes into potential themes; (4) reviewing themes and refining them to ensure coherence with both the coded data and the entire corpus; (5) defining and naming the themes; and (6) producing the final report. The process was supported by NVivo 12 qualitative data analysis software, which facilitated the management and organization of the emerging codes and themes.

This inductive analysis yielded three main thematic categories that structure the discussion of the review findings: 1) Dopamine as a signal for learning and motivational effort, which encompasses evidence regarding the role of dopamine in reward prediction, incentive salience attribution, and the mobilization of effort required for goal-directed actions; 2) Corticostriatal circuits in action selection

and persistence, which addresses the functional interaction between the ventral and dorsal striatum and prefrontal cortical regions in decision-making, action planning, and the maintenance of behavior in the face of costs and delays; and 3) From neurobiology to psychopathology: motivational dysregulation in depression and addiction, which synthesizes how hypoactivity and hyperactivity of reward and motivational circuits manifest clinically, focusing on anhedonia, apathy, and compulsive reward-seeking behavior.

Results and Discussion

Analysis of the 11 selected studies indicated that motivation to act toward a goal arises from complex, hierarchical interactions within the brain. The evidence reinforces the view that this process fundamentally depends on two interconnected systems: a chemical signaling system centered on dopamine and a network of brain structures that translates this signal into action. As the data demonstrate, dopamine does not function merely as a marker of pleasure; rather, it acts as an internal teaching signal, guiding learning about what is worth pursuing and adjusting the effort invested in these pursuits. However, this neurochemical learning process would be disorganized without the structural framework provided by circuits connecting deeper brain regions associated with desire to the prefrontal cortex, which is responsible for planning and persistence. This integration transforms an impulse into sustained, goal-directed behavior.

Understanding this dynamic extends beyond scientific inquiry. When the fine-tuned coordination between dopamine signaling and cortical circuit function is disrupted, the consequences can be clinically severe and manifest in opposite directions. On the one hand, an underactive system contributes to apathy and anhedonia, which are core symptoms of disorders such as major depressive disorder.

On the other hand, an overactive or “kidnapped” system leads to compulsive and pathological reward-seeking, as occurs in substance use disorders. Thus, the three lines of discussion structuring this section - the role of dopamine in motivational effort, the organization of brain circuits underlying action, and the clinical manifestations of their dysfunction - are not separate topics. Rather, there are complementary perspectives on the same overarching question: how the brain generates the drive to act and how this process can sometimes fail.

Dopamine as a Signal for Learning and Motivational Effort

Recent studies increasingly indicate that dopamine acts as an internal teaching signal, guiding learning about what is worth pursuing in the environment. When an action produces an outcome that is better than expected, a burst of dopamine release in the brain “marks” that behavioral sequence as successful, strengthening the neural connections that supported it. It is as though the brain were signaling, “This worked; remember to do it again.” Conversely, when an expected reward fails to materialize, dopaminergic activity decreases, indicating that the strategy



may need to be revised. This reward prediction error mechanism, originally proposed by Schultz, functions as a continuous updating system that helps optimize behavior in an ever-changing environment^{4,7}.

The reinforcement learning literature describes reward prediction error as the difference between an obtained outcome and the expected outcome. This signal can contribute to updating expectations and guiding choices. Contemporary reviews nevertheless emphasize that the relationship between dopaminergic signaling, learning, and behavior is more complex than a simple correspondence between dopamine and reward: the outcome depends on the context, the task, and the cognitive operations involved^{19,25,33}.

The distinction between wanting (motivation to pursue a reward) and liking (the hedonic response to consuming it) remains useful for organizing the literature, but it should not be interpreted as an absolute separation. Reviews on anhedonia indicate that pleasure, value assessment, anticipation, and learning are partially dissociable components, and that evidence of impairment in each component varies across studies and populations^{20,27,33}.

In experimental studies, alterations in dopaminergic transmission can modify the willingness to perform effortful actions without necessarily abolishing basic hedonic responses. For example, the review by Bailey and colleagues separately examined the contributions of D2 receptors to effort-based decisions and value-based decisions. Together with reviews on the neurobiology of effort, this literature suggests that dopamine should not be characterized simply as the “pleasure neurotransmitter” or, conversely, exclusively as the “seeking neurotransmitter”²²⁻²⁴.

Motivation can also be driven by abstract rewards and long-term goals, but the available evidence does not support attributing all such behaviors to a single circuit. Studies and reviews of effort-based decision-making describe the integrated involvement of the ventral striatum, prefrontal regions, and cingulate areas in estimating expected value, weighing costs, and preparing responses^{23,26,28}.

Thus, dopamine should be understood as a modulator of motivational and learning processes that operates within circuits involved in evaluating outcomes, costs, and opportunities. The capacity to initiate a task, persist in the face of obstacles, and adjust behavior cannot be reduced to a single neurotransmitter; rather, it depends on multiple systems and their interactions with the environment^{19,23,26}.

This understanding substantially changes how behavioral initiation is conceptualized. The energy required to begin a difficult task, the drive to get out of bed, and the willingness to persist despite an obstacle have neurochemical underpinnings that differ from the satisfaction experienced upon completion. A healthy dopaminergic system therefore acts as a spark that helps translate intention into action. When this system malfunctions, whether through reduced activity, as may occur in depression, or through hyperactivation focused on

a single goal, as may occur in addiction, the overall balance is disrupted. Recognizing that dopamine is primarily involved in motivation and effort, rather than simply pleasure, is therefore an important step toward developing more effective treatments for disorders that undermine the capacity to act.

Corticostriatal Circuits in Action Selection and Persistence

For motivation to translate into action, the evaluation of goal value must interact with systems responsible for action selection, planning, and execution. Neuroimaging reviews and studies of effort-based decision-making describe the involvement of prefrontal, striatal, and cingulate regions, with their respective contributions varying according to the type of choice and the cost of the required response. This literature supports a distributed network model rather than the assignment of a fixed causal sequence to specific brain regions based on isolated studies^{23,26,28}.

This integration of valuation and control provides a useful framework for understanding goal-directed behavior, although the strength of the evidence depends on the paradigm employed. Reviews of effort-based decision-making describe distinct contributions of prefrontal regions, the ventral striatum, and the anterior cingulate cortex to reward evaluation, choice monitoring, and preparation for effort. These findings support the involvement of multiple regions in action selection without demonstrating that any single connection functions as an on-off switch for motivation^{23,26,28}.

Maintaining a plan in the face of distractions and costs involves cognitive control, monitoring, and goal updating. The literature on motivation and fatigue proposes that control networks interact with dopaminergic systems to weigh expected benefits against the costs of effort. However, this does not establish that the dorsolateral prefrontal cortex alone is responsible for all forms of persistence or self-regulation²⁶.

The prefrontal cortex contributes to executive functions relevant to goal maintenance and response regulation, but goal-directed behavior depends on distributed networks. Studies of motivation and fatigue indicate that interactions among cognitive control, effort costs, and expected benefits may vary according to an individual's state and the demands of the task. Therefore, difficulties with persistence should not automatically be attributed to an isolated brain region^{26,28}.

Individual differences in motivation and persistence should be interpreted cautiously. Tasks used to assess effort-based decisions differ in the type of effort required, the nature of the reward, and the way task demands are communicated. This methodological heterogeneity complicates direct comparisons across studies and reinforces the need to describe the measures used precisely before relating performance to stable traits or brain connectivity²⁸.

These variations in corticostriatal circuits offer a neurobiological explanation for phenomena often attributed solely to “willpower.” The ability to persist with a difficult project, study for an examination rather than become



distracted, or save money for a future goal may be deeply rooted in the efficiency with which valuation regions, such as the ventral striatum, communicate with executive control regions, including the dorsolateral prefrontal cortex (DLPFC) and dorsal striatum. Recognizing this has important implications. On the one hand, it may help reduce the stigma associated with motivational difficulties by demonstrating that they can have a measurable biological substrate. On the other hand, it may inform the development of more targeted interventions, such as cognitive training designed to strengthen relevant functions or therapies intended to modulate neural activity, providing additional strategies to help individuals whose pursuit of goals is repeatedly disrupted by dysfunction in these circuits.

From Neurobiology to Psychopathology

In depression, anhedonia may involve alterations in reward anticipation, motivation to act, value assessment, and learning from outcomes. Systematic neuroimaging reviews describe alterations in striatal and frontocortical function, but the effects vary according to the component of reward processing assessed and the paradigm employed. Therefore, it is inappropriate to characterize all patients as exhibiting a uniform neural response or to attribute their symptoms to a single mechanism^{20,27,32,33}.

The relationship between depression, anhedonia, and effort was examined in a systematic review of 43 studies, which identified a reduced willingness to expend physical and cognitive effort for rewards among individuals with depression or anhedonia, with more consistent effects for physical effort. Neuroimaging literature also indicates differences between the anticipation and receipt of monetary and natural rewards. These findings support the separate assessment of motivational and hedonic components but do not establish that a specific frontostriatal disconnection is present in all depressive disorders^{21,32,33}.

In substance use disorders, reactivity to drug-associated cues is a widely used paradigm for investigating craving and motivational processes. A systematic review covering 25 years of fMRI studies identified hundreds of investigations involving nicotine, alcohol, cocaine, and other substances. However, it also highlighted methodological heterogeneity and the absence of cue-reactivity biomarkers approved for routine clinical use. Accordingly, brain activation in response to drug-related cues should be interpreted in the context of the experimental design rather than as isolated evidence of a specific dopaminergic mechanism^{30,31}.

Neuroimaging meta-analyses of craving identify the involvement of mesocorticolimbic networks and other brain regions, with variations according to the substance and sample characteristics. Recent methodological recommendations emphasize the importance of adequately reporting participant characteristics, tasks, craving assessments, and image acquisition and analysis procedures. Therefore, specific claims regarding dopamine release following chronic nicotine exposure or a concomitant reduction in responses to natural rewards should be made

Considered together, depression and addiction appear to represent two sides of the same coin: dysregulation of the mesocorticolimbic motivational system. Both can be conceptualized, in essence, as disorders of volition, affecting the capacity to direct behavior flexibly and adaptively. In depression, the system may be underactive, with a marked reduction in the perceived value of effort and a diminished capacity to generate motivation. In addiction, the system may become persistently biased toward a specific stimulus, resulting in excessive, rigid, and maladaptive motivation. Both conditions illustrate that healthy functioning depends not merely on the presence or absence of dopamine but on precise regulation and integrated communication between regions involved in value encoding, such as the ventral striatum, and those responsible for planning and controlling action, including the prefrontal cortex and dorsal striatum.

This unified understanding has important therapeutic implications. It suggests that interventions for depression should aim not only to increase dopaminergic tone but also to restore effective communication between the reward system and the prefrontal cortex, potentially through psychotherapies that promote action planning or through neuromodulation techniques. In addition, a dual strategy may be warranted: reducing the pathological salience attributed to drug-related cues while strengthening prefrontal cognitive control mechanisms to help individuals inhibit impulses and engage in alternative goals. Ultimately, the objective is not to suppress motivation but to restore its balance and redirect it toward adaptive goals.

Conclusion

This analysis provides a more comprehensive account of the processes that drive human behavior. It begins with a neurochemical signal associated with desire and opportunity, generated by dopamine in deeper brain regions, and culminates when this motivational signal interacts efficiently with the prefrontal cortex, which supports planning and control. Motivation is therefore not a simple on-off switch but a sequence of neural events that must be precisely coordinated. Clinical observations are consistent with this perspective. When this coordination fails, the consequences may include the behavioral disengagement observed in depression, in which effort no longer seems worthwhile, or the compulsive behavior characteristic of addiction, in which motivational drive becomes directed toward a single destructive goal. These conditions represent two sides of the same coin: disruption of the normal coordination between wanting and acting.

Understanding motivation as an interaction among different brain regions suggests new directions for treatment. Rather than focusing exclusively on adjusting the levels of a single neurotransmitter, strategies should aim to help these regions reconnect and function together more effectively. This may involve psychotherapies that actively train action planning and frustration tolerance or techniques such as neuromodulation that may help modify patterns of



brain communication. The therapeutic target thus shifts from a specific molecule to the functional integrity of an entire network. Nevertheless, much remains to be discovered. Further research is needed to determine why some individuals exhibit greater persistence, whereas others are more vulnerable to dysfunction in these circuits. Genetic factors, life experiences, and social context are likely to contribute to shaping this neural architecture. Longitudinal studies examining how these circuits change over time and in response to different therapies will be essential. A major future challenge will be to identify ways, potentially through brain imaging or other biomarkers, to predict which therapeutic approach is most likely to benefit each

individual, thereby enabling genuinely personalized treatment. Ultimately, neuroscience is demonstrating that the drive to act, whether to get out of bed, write a book, or overcome an addiction, has a complex and fascinating physical basis. Understanding this basis does not diminish the human experience; rather, it provides a more precise language for explaining the disruption of motivation and, more importantly, a clearer framework for developing interventions. This knowledge brings us closer to the goal of not merely treating symptoms but restoring the fundamental capacity to want, act, and pursue a meaningful path.

References

1. Salamone JD, Correa M. The mysterious motivational functions of mesolimbic dopamine. *Neuron*. 2012;76(3):470-85. <https://doi.org/10.1016/j.neuron.2012.10.021>
2. Berridge KC, Robinson TE. Liking, wanting, and the incentive-sensitization theory of addiction. *Am Psychol*. 2016;71(8):670-9. <https://doi.org/10.1037/amp0000059>
3. Pessoa L. On the relationship between emotion and cognition. *Nat Rev Neurosci*. 2008;9(2):148-58. <https://doi.org/10.1038/nrn2317>
4. Schultz W. Dopamine reward prediction-error signalling: a two-component response. *Nat Rev Neurosci*. 2016;17(3):183-95. <https://doi.org/10.1038/nrn.2015.26>
5. Wise RA. Dopamine, learning and motivation. *Nat Rev Neurosci*. 2004;5(6):483-94. <https://doi.org/10.1038/nrn1406>
6. Berridge KC. From prediction error to incentive salience: mesolimbic computation of reward motivation. *Eur J Neurosci*. 2012;35(7):1124-43. <https://doi.org/10.1111/j.1460-9568.2012.07990.x>
7. Schultz W. Neuronal reward and decision signals: from theories to data. *Physiol Rev*. 2015;95(3):853-951. <https://doi.org/10.1152/physrev.00023.2014>
8. Niv Y. Reinforcement learning in the brain. *J Math Psychol*. 2009;53(3):139-54. <https://doi.org/10.1016/j.jmp.2008.12.005>
9. Haber SN, Knutson B. The reward circuit: linking primate anatomy and human imaging. *Neuropsychopharmacology*. 2010;35(1):4-26. <https://doi.org/10.1038/npp.2009.129>
10. Balleine BW, O'Doherty JP. Human and rodent homologues in action control: corticostriatal determinants of goal-directed and habitual action. *Neuropsychopharmacology*. 2010;35(1):48-69. <https://doi.org/10.1038/npp.2009.131>
11. Der-Avakian A, Markou A. The neurobiology of anhedonia and other reward-related deficits. *Trends Neurosci*. 2012;35(1):68-77. <https://doi.org/10.1016/j.tins.2011.11.005>
12. Volkow ND, Wise RA, Baler R. The dopamine motive system: implications for drug and food addiction. *Nat Rev Neurosci*. 2017;18(11):741-52. <https://doi.org/10.1038/nrn.2017.130>
13. Botvinick M, Braver T. Motivation and cognitive control: from behavior to neural mechanism. *Annu Rev Psychol*. 2015;66:83-113. <https://doi.org/10.1146/annurev-psych-010814-015044>
14. Souza MT, Silva MD, Carvalho R. Revisão integrativa: o que é e como fazer. *Einstein (São Paulo)*. 2010;8(1):102-6. <https://doi.org/10.1590/s1679-45082010rw1134>
15. Whittemore R, Knafl K. The integrative review: updated methodology. *J Adv Nurs*. 2005;52(5):546-53. <https://doi.org/10.1111/j.1365-2648.2005.03621.x>
16. Botelho LLR, Cunha CCA, Macedo M. O método da revisão integrativa nos estudos organizacionais. *Gest Soc*. 2011;5(11):121-36. <https://doi.org/10.21171/ges.v5i11.1220>
17. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;372:n71. <https://doi.org/10.1136/bmj.n71>
18. Braun V, Clarke V. Using thematic analysis in psychology. *Qual Res Psychol*. 2006;3(2):77-101. <https://doi.org/10.1191/1478088706qp063oa>
19. Papalini S, Beckers T, Vervliet B. Dopamine: from prediction error to psychotherapy. *Transl Psychiatry*. 2020;10:164. <https://doi.org/10.1038/s41398-020-0814-x>
20. Borsini A, St John Wallis A, Zunszain P, Pariante CM, Kempton MJ. Characterizing anhedonia: a systematic review of neuroimaging across the subtypes of reward processing deficits in depression. *Cogn Affect Behav Neurosci*. 2020;20(4):816-41. <https://doi.org/10.3758/s13415-020-00804-6>
21. Horne SJ, Topp ET, Quigley L. Depression and the willingness to expend cognitive and physical effort for rewards: a systematic review. *Clin Psychol Rev*. 2021;88:102065. <https://doi.org/10.1016/j.cpr.2021.102065>
22. Bailey MR, Chun E, Schipani E, Balsam PD, Simpson EH. Dissociating the effects of dopamine D2 receptors on effort-based versus value-based decision making using a novel behavioral approach. *Behav Neurosci*. 2020;134(2):101-18. <https://doi.org/10.1037/bne0000361>
23. Salamone JD, Correa M. The neurobiology of motivational aspects of motivation: exertion of effort, effort-based decision making, and the role of dopamine. *Annu Rev Psychol*. 2024;75:1-32. <https://doi.org/10.1146/annurev-psych-020223-012208>



24. Salamone JD, Ecevitoglu A, Carratala-Ros C, Presby RE, Edelstein GA, Fleeher R, et al. Complexities and paradoxes in understanding the role of dopamine in incentive motivation and instrumental action: exertion of effort vs. anhedonia. *Brain Res Bull.* 2022;182:57-66. <https://doi.org/10.1016/j.brainresbull.2022.01.019>
25. Corlett PR, Mollick JA, Kober H. Meta-analysis of human prediction error for incentives, perception, cognition, and action. *Neuropsychopharmacology.* 2022;47(7):1339-49. <https://doi.org/10.1038/s41386-021-01264-3>
26. Kok A. Cognitive control, motivation and fatigue: a cognitive neuroscience perspective. *Brain Cogn.* 2022;160:105880. <https://doi.org/10.1016/j.bandc.2022.105880>
27. Wang S, et al. Anhedonia as a central factor in depression: neural mechanisms revealed from preclinical to clinical evidence. *Prog Neuropsychopharmacol Biol Psychiatry.* 2021;110:110289. <https://doi.org/10.1016/j.pnpbp.2021.110289>
28. Liang S, Wu Y, Li H, Greenshaw AJ, Li T. Anhedonia in depression and schizophrenia: brain reward and aversion circuits. *Neuropsychiatr Dis Treat.* 2022;18:1385-96. <https://doi.org/10.2147/NDT.S367839>
29. Ekhtiari H, Zare-Bidoky M, Sangchooli A, et al. A methodological checklist for fMRI drug cue reactivity studies: development and expert consensus. *Nat Protoc.* 2022;17(3):567-95. <https://doi.org/10.1038/s41596-021-00649-4>
30. Addiction Cue-Reactivity Initiative (ACRI) Network, et al. Parameter space and potential for biomarker development in 25 years of fMRI drug cue reactivity: a systematic review. *JAMA Psychiatry.* 2024;81(4):414-25. <https://doi.org/10.1001/jamapsychiatry.2023.5483>
31. Pollard AA, Hauson AO, Lackey NS, et al. Functional neuroanatomy of craving in heroin use disorder: voxel-based meta-analysis of functional magnetic resonance imaging (fMRI) drug cue reactivity studies. *Am J Drug Alcohol Abuse.* 2023;49(4):418-30. <https://doi.org/10.1080/00952990.2023.2172423>
32. Bore MC, Liu X, Gan X, et al. Distinct neurofunctional alterations during motivational and hedonic processing of natural and monetary rewards in depression: a neuroimaging meta-analysis. *Psychol Med.* 2024;54(4):639-51. <https://doi.org/10.1017/S0033291723003410>
33. Kieslich K, Valton V, Roiser JP. Pleasure, reward value, prediction error and anhedonia. *Curr Top Behav Neurosci.* 2022;58:281-304. https://doi.org/10.1007/7854_2021_295

